

# Extreme star formation in starburst galaxies



# Extreme star formation in starburst galaxies

## Proefschrift

ter verkrijging van  
de graad van Doctor aan de Universiteit Leiden,  
op gezag van Rector Magnificus prof. mr. dr. P. F. van der Heijden,  
volgens besluit van het College voor Promoties  
te verdedigen op woensdag 28 november 2007  
klokke 16.15 uur

door

Leonie Snijders

geboren te Beverwijk  
in 1979

Promotiecommissie

Promotor: Prof. dr. M. Franx

Co-promotor: Dr. P. P. van der Werf

Referent: Prof. R. C. Kennicutt, Jr. (University of Cambridge, UK)

Overige leden: Dr. B. R. Brandl  
Prof. dr. E. F. van Dishoeck  
Prof. dr. F. P. Israel  
Dr. L. J. Kewley (University of Hawaiï, Honolulu, USA)  
Prof. dr. K. H. Kuijken  
Prof. dr. P. T. de Zeeuw



Cover:  
*Liten hägring (Minor Mirage)*,  
Henrik Samuelsson, Sweden 2003.  
Courtesy Galerie Laurent Godin, Paris.

# Table of contents

	Page
<b>Chapter 1. Introduction</b>	<b>1</b>
1.1 Starburst galaxies . . . . .	1
1.2 Observational characteristics of starburst galaxies . . . . .	2
1.3 This thesis . . . . .	3
1.4 Objects . . . . .	4
1.4.1 The Antennae . . . . .	4
1.4.2 M83 . . . . .	5
1.5 Observations . . . . .	6
1.5.1 Ground-based . . . . .	6
1.5.2 Space-based . . . . .	7
<b>Chapter 2. Mid-infrared diagnostics of starburst galaxies</b>	<b>11</b>
2.1 Introduction . . . . .	12
<b>Part I: Modeling emission line spectra of embedded star clusters</b>	<b>14</b>
2.2 Modeling the emission line spectra . . . . .	15
2.2.1 Spectral features in the wavelength range of the atmospheric windows . . . . .	15
2.2.2 Stellar population synthesis model . . . . .	16
2.2.3 Photoionization model . . . . .	19
2.2.4 Parameter space covered by the model grid . . . . .	20
2.3 Model results . . . . .	21
2.3.1 Age evolution of line ratios: mid-infrared . . . . .	21
2.3.2 Age evolution of line ratios: near-infrared . . . . .	26
2.4 Comparison with existing mid-infrared models . . . . .	28
2.5 Diagnostic diagrams . . . . .	30
2.6 Comparison with published data . . . . .	33
<b>Part II: A test case for model applications</b>	<b>34</b>
2.7 Stellar clusters in the overlap region of the Antennae . . . . .	35
2.7.1 Observations, data reduction and results . . . . .	35
2.8 Discussion . . . . .	39
2.8.1 Modeling infrared line emission of young star-forming regions . . . . .	39
2.8.2 Dense star-forming regions in the Antennae . . . . .	40
2.8.3 Evidence for a dense, clumpy ISM . . . . .	42
2.9 Conclusions . . . . .	46
<b>Chapter 3. Ground-based VLT mid-infrared observations of NGC 4038/4039</b>	<b>49</b>
3.1 Introduction . . . . .	50
3.2 Observations and data reduction . . . . .	50

3.3 Results . . . . .	53
3.4 PAH emission . . . . .	55
3.5 Future work . . . . .	57
<b>Chapter 4. Ground-based VLT near-infrared spectroscopy of NGC 4038/4039</b>	<b>59</b>
4.1 Introduction . . . . .	60
4.2 Observations and data reduction . . . . .	61
4.3 Results . . . . .	62
4.4 Cluster properties . . . . .	63
4.4.1 Extinction . . . . .	66
4.4.2 Cluster ages . . . . .	66
4.4.3 $T_{\text{eff}}$ of the stellar radiation field . . . . .	68
4.4.4 Cluster masses . . . . .	70
4.5 The excitation mechanism of molecular hydrogen . . . . .	72
4.6 Conclusions . . . . .	75
<b>Chapter 5. Space-based Spitzer mid-infrared spectroscopy of NGC 4038/4039</b>	<b>81</b>
5.1 Introduction . . . . .	82
5.2 Observations and data reduction . . . . .	85
5.2.1 ‘Hires’ observations . . . . .	85
5.2.2 ‘Lores’ observations . . . . .	87
5.3 Results and methods . . . . .	91
5.3.1 Continuum fluxes . . . . .	91
5.3.2 Polycyclic aromatic hydrocarbons (PAHs) . . . . .	93
5.3.3 Fine-structure and hydrogen emission lines . . . . .	95
5.3.4 Deriving $H_2$ temperatures and masses . . . . .	96
5.3.5 Extinction estimates . . . . .	100
5.4 Discussion . . . . .	101
5.4.1 The nuclei of NGC 4038 and NGC 4039 . . . . .	102
5.4.2 The properties of individual SSCs . . . . .	104
5.4.3 Cluster ages in comparison . . . . .	106
5.4.4 The physical conditions across the Antennae . . . . .	107
5.4.5 Dust temperature and radiation field . . . . .	109
5.4.6 Star formation rates for the individual star-forming regions . . . . .	109
5.4.7 PAHs – Strengths and variations . . . . .	110
5.4.8 The strength of the $H_2$ emission . . . . .	113
5.4.9 $H_2$ temperatures and masses . . . . .	115
5.4.10 $H_2$ excitation mechanisms . . . . .	118
5.5 Summary . . . . .	119
<b>Chapter 6. PAH emission in the nucleus of M83</b>	<b>123</b>
6.1 Introduction . . . . .	124
6.2 Observations and data reduction . . . . .	125
6.2.1 VISIR narrowband imaging . . . . .	125
6.2.2 VISIR spectroscopy . . . . .	126

6.2.3 SINFONI data . . . . .	128
6.3 Results . . . . .	129
6.4 Diffuse versus compact PAH emission . . . . .	132
6.5 Properties of the star-forming regions . . . . .	136
6.5.1 The central star formation complex . . . . .	137
6.5.2 Source 8 . . . . .	139
6.6 Conclusions . . . . .	140
<b>Chapter 7. Summary &amp; Outlook</b>	<b>143</b>
7.1 Models . . . . .	143
7.2 Highlights . . . . .	144
7.2.1 Properties of the stellar populations and the surrounding ISM . . .	144
7.2.2 Diffuse PAH emission . . . . .	146
7.3 Outlook . . . . .	147
<b>Nederlandse Samenvatting</b>	<b>149</b>
<b>Curriculum Vitae</b>	<b>157</b>
<b>Nwoord</b>	<b>158</b>

### **Alheimsvíðáttan**

Sá eg í ungum  
æskublóma  
stjörnur úr himin-  
straumum rísa,  
þúsund alda  
aðþreyta skeið  
heiðfagran gegnum  
himinbláma.

(...)

Gránar í geimi,  
geysa ég um himin  
þokuþungaðan  
þjótandi fram;  
dunar mér á baki  
dökknaðra sóla  
flugniður allra,  
sem fossa deyjandi.

### **The Vastness of the Universe**

I watched the stars  
in the womb of youth  
rise from the still  
streams of heaven,  
eager to make  
their million year  
race through the thin  
ethereal blue.

(...)

As I dart on  
through dim greyness,  
I encounter clouds  
of cosmic dust.  
Behind me I hear,  
hushed in distance,  
dark cataracts  
of dying suns.

**Jónas Hallgrímsson, Iceland, 1837**